

Deciphering the Benson-Bassham-Calvin Cycle

1. A technological revolution for deciphering the cycle



Figure 1. Discoverers of the photosynthetic carbon reduction cycle (from left to right): James A. Bassham, Andrew A. Benson and Melvin Calvin. Photo presented to Karl Biel by Andrew A. Benson (September 9, 1988). [Source: Reprinted with permission of Karl Biel]

Around 1945, the american chemists A.A. Benson, J.A. Bassham and M. Calvin [\[1\]](#) (Figure 1) tackled the task of identifying the first carbon-containing product of photosynthesis that had escaped all scientific investigation. After the World War II, these researchers had just acquired two new important technological allies in Berkeley, California:

- a radioactive tracer, ^{14}C , which had just been isolated by S. Ruben and M. Kamen in 1941 [\[2\]](#),
- new techniques for the chromatographic separation of the carbon-containing compounds from metabolism, developed a few years earlier in England by R. Martin and A. Syngé. [\[3\]](#)

These two tools will allow Dr. Calvin and his co-workers to identify the path taken by carbon during the various biochemical reactions for assimilation of the photosynthetic carbon.



Figure 2. Diagram of the apparatus used to monitor the incorporation of photosynthetic carbon into the products of photosynthesis . In the centre: device used for the study of CO₂ fixation by unicellular algae. [Source Photo © Ernest O. Lawrence, Berkeley National Laboratory, Photography and Digital Imaging Services: previously unpublished, *Biochemistry & Molecular Biology of Plants*. B.B. Buchanan, W. Gruissem & R.L. Jones. American Society of Plant Physiologists, Rockville, MD. 2000 p. 610]. Right: Lollipop given to Pr. Roland Douce by Andrew A. Benson and kept at the Musée des Confluences (Lyon, France).

These first experiments were not carried out on leaves but on unicellular algae (*Chlorella* or *Senedesmus*), which were easier to use. The experimental protocol is as follows:

1. The unicellular algae are placed in a suspension medium that flows at a constant rate from a tank through a transparent illuminated coil.
2. When photosynthesis is stationary, under illumination at a given temperature, sodium bicarbonate (Na¹⁴CO₃) is introduced by a syringe into the suspension medium.
3. Incubation in the presence of ¹⁴C is limited to the time the suspension continues to descend through the coil after injection (a few seconds).
4. At the end, the plant material is fixed in boiling methanol (Figure 2). [4]
5. The constituents present in the extract are then separated by two-dimensional chromatography.
6. The chromatogram is then X-rayed in the dark. The black spots on the radio-chromatogram indicate the presence of the radioactive compounds that were formed during the ¹⁴C labelling of the algae.
7. In order to identify these compounds, radiochromatograms are compared with chromatograms coloured by different chemical reagents specific to organic acids, amino acids, sugars and phosphorylated compounds (Figure 3).




Figure 3. Auto-radiograms of ¹⁴C-labelled photosynthetic compounds after separation by two-dimensional chromatography [Source: Scheme R Prat, in Morot-Gaudry, Dunod, 2009]

2. The Benson-Bassham-Calvin Cycle




Figure 4. Schematic representation of the Benson-Bassham-Calvin Cycle. The molecules are shown only with their carbons and phosphate groups.

All the molecules identified with the methods described above were organized in a cycle comprising 3 phases. Figure 4 schematically depicts the Benson-Bassham-Calvin Cycle.

Phase 1 - Carbon Dioxide (CO₂) fixation

$3 \text{ RuBP (C5 molecule)} + 3 \text{ CO}_2 \text{ (C1 molecule)} \rightarrow 6 \text{ 3-PGA (C3 molecule)}$ (catalyzed by the Rubisco)

Phase 2 - Reduction of phosphoglyceric acid to triose-phosphate

The 3-phosphoglyceric acid (3-PGA) is reduced in two steps to glyceraldehyde 3-phosphate (G3P). A molecule with 3 carbons and a phosphate group, G3P is a triose phosphate. Both reactions consume NADPH₂ and part of the ATP produced by thylacoids during the photochemical reactions of photosynthesis (see [Shedding light on Photosynthesis](#)):

$6 \text{ 3-PGA (C3 molecule)} + 6 \text{ ATP} \rightleftharpoons 6 \text{ 1,3-bisPGA (C3 molecule)} + 6 \text{ ADP}$

$6 \text{ 1,3-bisPGA (C3 molecule)} + 6 \text{ NADPH} + 6 \text{ H}^+ \rightleftharpoons 6 \text{ G3P (C3 molecule)} + 6 \text{ NADP}^+ + 6 \text{ Pi}$

Phase 3 - Regeneration of the CO₂ acceptor

The role of the cycle is to regenerate the RuBP. The set of reactions in phase 3 consists of reconvertng molecules with 3 carbon atoms into molecules with 5 carbon atoms in order to be able to reset the cycle. This phase consumes very large quantities of glyceraldehyde 3-phosphate (5 molecules out of 6 formed). Phosphorylated C₆, C₄, C₇ and finally C₅ backbones are made (see Figure 4), as can be represented schematically as follows:

$3\text{C}3\text{P} + 3\text{C}3\text{P} \rightarrow \text{C}6\text{-diP} \rightarrow \text{C}6\text{-P} + \text{Pi}$

$\text{C}6\text{-P} + 3\text{C}3\text{P} \rightarrow \text{C}4\text{P} + \text{C}5\text{P}$

$\text{C}4\text{P} + 3\text{C}3\text{P} \rightarrow \text{C}7\text{-diP} \rightarrow \text{C}7\text{-P} + \text{Pi}$

$\text{C}7\text{P} + 3\text{C}3\text{P} \rightarrow 2 \text{ C}5\text{P}$

Only the last stage of this phase consumes ATP. The overall balance sheet can be summarized as follows:

$5 \text{ G3P (C3 molecule)} \rightarrow 3 \text{ ribulose-5-P (C5 molecule)} + 2 \text{ Pi}$

$3 \text{ ribulose-5-P (C5 molecule)} + 3 \text{ ATP} \rightleftharpoons 3 \text{ RuBP (C5 molecule)} + 3 \text{ ADP}$

To summarize, only one of the 6 triose phosphate molecules produced by this cycle will be used for the production of sugars and, more broadly, other metabolites that are the basis of biomass creation (Figure 4). It will be used for the biosynthesis of starch,

amino acids or lipids in the chloroplast or exported out of the chloroplast and transformed into sucrose by the enzymes of the cytoplasm (see Focus [Sucrose or Starch?](#)).

Notes and References

Cover image. Structures of some molecules of the Benson-Bassham-Calvin Cycle (Public domain)

[1] Andrew Alm Benson (1917-2015), James Alan Bassham (1922-2012) & Melvin Calvin (1911-1997), are American chemists whose contributions to the discovery of the carbon cycle in plants are considerable. Benson, a specialist in carbon compounds and sugars, is credited in particular with the discovery of ribulose 1,5-bisphosphate on which the carbon of CO₂ is fixed. Calvin was awarded the 1961 Nobel Prize in Chemistry for his work on the assimilation of carbon dioxide by plants.

[2] Martin Kamen (1913-2002) and Samuel Ruben (1913-1943) are two American chemists, co-discoverers of carbon-14 (¹⁴C). With a half-life of 5730 years, ¹⁴C can be very easily used in metabolic experiments. It is one of the three naturally abundant isotopes (along with ¹²C and ¹³C), but the only radioactive one.

[3] Richard Laurence Millington Synge (1914-1994), and Archer John Porter Martin (1910-2002) are both English chemists. They were co-winners of the 1952 Nobel Prize in Chemistry for the invention of "partition chromatography".

[4] One of the devices -offered by Andrew Benson to Roland Douce- is kept at the Musée des Confluences in Lyon (France).

More:

- Farineau J. & Morot-Gaudry F., 2011, La Photosynthèse, Quae, ISBN 978-2-7592-0903-3
- Morot-Gaudry F., Moreau F., Prat R., Maurel C. & Sentenac H. (2017) Biologie végétale : Nutrition et métabolisme - 3e édition, Dunod
- Video: [A conversation between Andrew Benson and Professor Bob Buchanan on the discovery of the Benson-Bassham-Calvin Cycle](#)

L'Encyclopédie de l'environnement est publiée par l'Université Grenoble Alpes.

Les articles de l'Encyclopédie de l'environnement sont mis à disposition selon les termes de la licence Creative Commons Attribution - Pas d'Utilisation Commerciale - Pas de Modification 4.0 International.
