

What can be done to restore dispersal ?

1. Restoring dispersal by sowing native trees



Figure 1: Direct sowing of large-seeded trees in December 2021 in one of the ECOFORRUN project plots. This project aims to restore tropical rainforest by combining invasive plant control with direct sowing. These 9 species are: 1, *Strongylodon lucens* (Fabaceae), 2, *Hyophorbe indica* (Arecaceae), 3, *Mimusops balata* (Sapotaceae), 4, *Hernandia mascarenensis* (Hernandiaceae), 5, *Diospyros borbonica* (Ebenaceae), 6, *Terminalia bentzoe* (Combretaceae), 7, *Pandanus purpurascens* (Pandanaceae), 8, *Ochrosia borbonica* (Apocynaceae), 9, *Cassine orientalis* (Celastraceae). [Source: photo © Alexis Gorissen, reproduced with permission of the author]

In Reunion Island, experimental sowings have shown that it is possible to restore the recruitment of various woody plants with large fleshy fruits on historical lava flows (see Figure 9, article [How vertebrate extinctions threaten tropical forests](#)). These experiments also tested the impact of other limitations such as seed predation by rats or competition with invasive plants. These experiments showed that the loss of dispersal due to the disappearance of frugivores is the main factor explaining such a depletion of native woody diversity [\[1\]](#).

Based on these encouraging results, we decided to continue the experiments in tropical rainforests (Figure 1).

The success of direct sowing opens new perspectives for ecological restoration because it is easier to implement than traditional sapling plantations. The latter require very heavy logistics to be deployed on a large scale. As already shown elsewhere [\[2\]](#), direct sowing would allow sowing of remote high-stake areas such as the volcano caldera, which has the largest lowland protected area in the Mascarene Islands (see Figure 4, article [How vertebrate extinctions are threatening tropical forests](#)). It is even more important to maintain the most iconic rainforest tree species in this area to maintain attractive ecosystems in anticipation of the return of large frugivores.

2. Restoring dispersal by rewilding ecosystems

Direct sowing is not intended to be an effective substitute for large frugivores. Sowing can only be undertaken in limited spatial and temporal scales compared to the dispersal capabilities of large fauna. Nothing will ever replace large numbers of foraging frugivores with long ranges. However, while direct sowings could be easily deployed by conservationists, *rewilding* raises a number of social-ecological issues that make it more difficult to implement.

2.1. Re-introduction of terrestrial tortoises



Figure 2. Giant Seychelles tortoise (*Aldabrachelys gigantea*) [Source: photo © [YotcmdrDerivative by User:MathKnight, CC BY 4.0, via Wikimedia Commons]

The exotic land tortoises *Aldabrachelys gigantea* (Figure 2) and *Astrochelys radiata* are considered to be good functional analogues of the extinct *Cylindraspis* spp. tortoises in the Mascarenes [3]. They have been successfully used in various ecological restoration projects in the archipelago. However, it is easier to rewild the islets of Mauritius with these tortoises than the native habitats of Reunion Island: these islets are isolated nature reserves, while the problems of securing the animals are paramount in Reunion Island where poaching is to be feared.

Nevertheless, the recent reintroduction of land tortoises to the *Ebony Forest* reserve [4] on the mainland of Mauritius shows that a similar initiative could see the light of day in Reunion. While such a project should not be expected to have an ecosystem-wide impact (at first), a reserve with land tortoises could not only be an exciting experiment in ecological restoration, but also have an important educational purpose. It would help to the public aware of the essential role large vertebrates play in native tropical forests.

2.2 The case of the black flying fox



Figure 3. Dormitory of the black dogfish (*Pteropus niger*) in eastern Reunion Island. The black dogfish recolonized Reunion Island from Mauritius in the early 2000s about two centuries after it had disappeared. [Source: photo © Gildas Monnier - GCOI, reproduced with the author's permission]

Figure 3. Roost of the Mauritian flying fox (*Pteropus niger*) in eastern Reunion Island. The Mauritian flying fox recolonized Reunion Island from Mauritius in the early 2000s about two centuries after it had disappeared. [Source: photo © Gildas Monnier – GCOI, reproduced with the author's permission]

The return of the Mauritian flying fox to Reunion after a cyclone is a remarkable opportunity for forest ecosystems (Figure 3). Their still very modest population is currently growing and it is to be hoped that this bat, once highly valued for its meat, will not be victim of poachers. However, the return of fruit bats to Reunion sland raises two main concerns for some conservationists.

- The risk of new plant invasions in an already severely impacted island;
- The risk of conflict of interest with tree farmers.

The risk of seeing the bats dispersing exotic plants exists. However, these potential negative effects will certainly be counterbalanced by the restoration of mutualistic pollination and dispersal interactions that have been extinct for centuries [5]. In particular, their ability to disperse seeds over long distances will restore gene flow for many native species that have been locked in their forested areas for over 150 years. To see the positive impact of Mauritian flying fox, one need only observe the remarkable regeneration of most native plants at Brise Fer despite the existence of many invasive exotic plants on the landscape scale in this massif (see Section 3.2 [How vertebrate extinctions threaten tropical forests](#)).

On Reunion Island, lowland native forests rich in large fleshy-fruited trees are often highly fragmented and in close proximity to cultivated areas. With the growth of the fruit bat population, the potential conflict of interest with arborists should be anticipated. The challenge is great as shown by the situation in Mauritius where this species, which plays a key role in the native forests, has been decimated in the last decade. Therefore, local conservationists should ensure that the conditions for the acceptance of this emblematic species are created and include an important socio-ecological component in future research. This work can only facilitate other reintroduction projects, such as the one concerning the Mascarene parakeet, which is being actively considered by several key conservation stakeholders in Reunion Island.

Notes

Cover image. Copy of one of the images in figure 7 of the main article.

[1] Albert S. (2020) Rupture des interactions mutualistes plantes à fruits charnus-vertébrés frugivores, et conséquences sur la régénération des forêts tropicales dans les Mascareignes, Doctoral dissertation, Université de La Réunion. [On line]. Available at: <https://tel.archives-ouvertes.fr/tel-03126708>

[2] Shaw N. *et al.* (2020) Seed use in the field: Delivering seeds for restoration success, *Restor. Ecol.*, 28, S3, S276-S285, <https://doi.org/10.1111/rec.13210>.

[3] Griffiths C. J., Hansen D. M., Jones C. G., Zuñel N. & Harris S. (2011) Resurrecting extinct interactions with extant substitutes, *Curr. Biol.* 21(9), 762-765 Available at: <http://www.sciencedirect.com/science/article/pii/S0960982211003460>

[4] Ebony forest reserve turtles: <https://www.ebonyforest.com/facilities/tortoises/>

[5] Florens F. B. V. *et al.* (2017) Disproportionately large ecological role of a recently mass-cultivated flying fox in native forests of an oceanic island, *J. Nat. Conserv.* 40, 85-93, <https://doi.org/10.1016/j.jnc.2017.10.002>

L'Encyclopédie de l'environnement est publiée par l'Université Grenoble Alpes.

Les articles de l'Encyclopédie de l'environnement sont mis à disposition selon les termes de la licence Creative Commons Attribution - Pas d'Utilisation Commerciale - Pas de Modification 4.0 International.
