

## Thermogenesis and pollination in Araceae

## 1. Respiratory crisis in Araceae



*Figure 1. Photo of an Arum maculatum inflorescence. The insert on the right shows the interior of the floral chamber observed after a longitudinal section. [Diagram: author's montage from photos © Laurent Francini, reproduced with the permission of the author]* 

In Araceae, during the development of inflorescences (called spadices), a strong increase in respiration (respiratory crisis) is observed, coupled with a strong increase in temperature. This phenomenon of overheating, called thermogenesis, was discovered by Lamarck in 1778 and the relationship between an increase in respiratory oxygen consumption and heat emission was demonstrated by Garreau in 1851 [1]. Figure 1 shows a plant of *Arum maculatum* with its flowers hidden in the floral chamber topped by a sterile club. This inflorescence is hidden inside a particular leaf, called a spathe, which rolls up on itself. A cut in the wall of the floral chamber makes it possible to distinguish the various types of grouped flowers and their organization. The inflorescence grows steadily through successive developmental stages (denoted **a** then  $\alpha$  to  $\zeta$ ) described by James and Beevers [2] (Figure 2) [3].



withered spathe and club, greenish ovules



The respiratory crisis of the spadix and, more specifically, the club, commonly occurs shortly after the spathe opens, as early as the  $\delta$  stage. It uses starch stored during development and lasts for a few hours during which the respiratory intensity and temperature of the spadix follow a parallel evolution [4] (Figure 3). The temperature of the spadix can thus reach 30°C and be nearly 20°C above air temperature.

## 2. Cyanide insensitive respiration coupled with heat production

The cyanide resistance of Araceae respiration, which is particularly important at the time of respiratory crisis, was observed as early as the 1950s [2]. Isolated spadix mitochondria exhibit electron transport using both the classical cyanide-sensitive cytochromic pathway terminated by the <u>cytochrome oxidase</u> complex and a cyanide-resistant, non-phosphorylating, non-cytochromic pathway, the latter of which is particularly active during the respiratory crisis [5].



Figure 3. Changes in respiratory intensity and spadix temperature at the time of respiratory crisis [Source: Author's diagram, adapted from ap Rees et al. [4]).

Figure 3 shows the evolution of mitochondrial oxidation of malate (organic acid at the origin of NADH reoxidized by the respiratory chain, see Figure 10 of the main article) and associated phosphorylation during the developmental stages of *Arum maculatum* spadix [6]. Malate oxidation peaks at the time of the respiratory crisis, which corresponds to the increase in

respiration. At the same time, the efficiency of oxidative phosphorylation (P/O), which corresponds to ATP production, already low in the early stages of development, decreases sharply during the respiratory crisis, indicating a significant involvement of the non-phosphorylating electron transport pathway [6]. The involvement of AOX, the terminal oxidase of this pathway, was subsequently demonstrated [7].

The low level of oxidative phosphorylation reflects the release of excess energy from the high level of respiration in the form of heat.

## 3. Heat production and pollination in Arum



*Figure 4. Evolution, over developmental stages, of malate oxidation and oxidative phosphorylation in mitochondria of Arum maculatum clubs (Author's diagram, adapted from Chauveau and Lance [6]).* 

Female *Arum maculatum* flowers become functional (i.e., susceptible to pollination) when the inflorescence begins to open. Within the spadix, thermogenesis has started: it promotes (*i*) the activity of enzymes catalyzing the synthesis of volatile amino compounds and (*ii*) the emission of these molecules in the vicinity of Arum plants. Their odour of excrement and urine attracts pollinating insects.

Pollination of the *Arum maculatum* flower is carried out by small 2-3 mm midges known as psychodes (*Psychoda phalaenoides*) [8] and also nicknamed "*sink midges*" or "*butterfly flies*" (Figure 5). These insects are very common in damp places (e.g., sewage pipes).



Figure 5. The "most read in the burrows" newspaper, The Hoot, described in detail the process that takes place in the heart of the inflorescence of the wild arum and that attracts a species of insect, Psychoda phalaenoides (or Psykoda), thus involuntarily participating in the pollination of the Arum. [Source: Reproduced from issue 26 (1985) of "la Hulotte" (DR)]

Moreover, the high temperature (15 to 20°C above room temperature) found within the spathe is very suitable for this small insect. *Psychoda* lands on the cone, slides, because the walls are very smooth, and passes without hindrance through the sterile

flowers. These flowers are so made that they trap *Psychoda* in the floral chamber: filaments pointing downwards (see Figure 1) prevent the insect from going up and coming out. It is then trapped in the lower part of the inflorescence (Figure 5) [9]. Many other *Psychoda* insects will join it and will also remain trapped: the Arum inflorescence is a real insect trap (Figure 6) [10] !

But things end up working out for *Psychoda*: after 2 days, the male flowers mature, open and let out pollen which sprinkles the midges, the sterile flowers wither and open the floral chamber... The imprisoned midges can then escape... They fly to other Arums which they will then be able to fertilize, in return for a new episode of imprisonment!



*Figure 6. During the respiratory crisis, many pollinating insects (Psychoda phalaenoides midge) are held captive in the floral chamber (the latter has been cut out to reveal its contents). [Photo © Laurent Francini, with permission of the author]* 

*Arum maculatum* and, more globally, the Araceae, are not the only species capable of thermogenesis and a recent article proposes a general model of thermoregulation which gives a central role to AOX [11].

Cover image. Arum Maculatum. Royalty free.

**Acknowledgements.** We are very grateful to Mr. Laurent Francini (Association La Chanterelle de Ville-la-Grand, Haute-Savoie) for the pictures which allowed to illustrate this focus on the pollination of Arum maculatum by the midge Psychoda phalaenoides.

[1] L. Garreau, 1851. De la respiration chez les plantes, Mémoire sur les relations qui existent entre l'oxygène, consommé par le spadice de l'*Arum italicum* en état de paroxysme, et la chaleur qui se produit, *Ann. Sci. Nat.* Ser. III, 15, 5-36. (in French)

[2] W.O. James & H. Beevers, 1950. The respiration of *Arum spadix*. A rapid respiration, resistant to cyanide, *New Phytol*. 49, 353-374. https://doi.org/10.1111/ j.1469-8137.1950.tb05160.x.

[3] Lance C 1972. Respiration of *Arum maculatum* during inflorescence development. *Annals of Natural Sciences*. *Botany* <sup>12th</sup> series, Volume XIII, 477-495.

[4] ap Rees T, Wright BW & Fuller WA (1977). Measurements of starch breakdown as estimates of glycolysis by the spadix of *Arum maculatum* L. *Planta*, 134, 53- 56. https://doi.org/10.1007/BF00390094.

[5] James WO, & Elliott DC (1955). Cyanide-resistant mitochondria from the spadix of an arum. *Nature*, 175, 89. 10.1038/175089a0

[6] Chauveau M & Lance C (1982). Respiration and thermogenesis in Araceae. *Bulletin de la Société Botanique de France. Actualités Botaniques*. 129, 123-134. <u>https://doi.org/10.1080/01811789.1982.10826566</u> (in French)

[7] Grant NM, Miller RE, Watling JR & Robinson SA (2008). Synchronicity of thermogenic activity, alternative pathway respiratory flux, AOX protein content, and carbohydrates in receptacle tissues of sacred lotus during floral development. *Journal of Experimental Botany*, 59, 705-714.

[8] https://www.francini-mycologie.fr/BOTANIQUE/Arum\_maculatum.html

[9] Libérez Psikoda ! Journal La Hulotte N° 26, January 1, 1985; <u>https://www.lahulotte.fr/</u> ((in French)

[10] https://www.francini-mycologie.fr/BOTANIQUE/Arum\_maculatum.html

[11] Wagner AM, Krab K, Wagner MJ & Moore AL (2008). Regulation of thermogenesis in flowering Araceae: the role of the alternative oxidase. *Biochimica Biophysica Acta*, 1777, 993-1000. <u>https://doi.org/10.1016/j.bbabio.2008.04.001</u>

L'Encyclopédie de l'environnement est publiée par l'Université Grenoble Alpes.

Les articles de l'Encyclopédie de l'environnement sont mis à disposition selon les termes de la licence Creative Commons Attribution - Pas d'Utilisation Commerciale - Pas de Modification 4.0 International.