



Plasmodium falciparum, a parasite derived from an algae

[Eric Maréchal](#), CNRS Research Director, Laboratory of Cellular & Plant Physiology, Université Grenoble Alpes.

A. First step: a primary endosymbiosis at the origin of red algae



B. Second step: a secondary endosymbiosis at the origin of Apicomplexa parasites



More than two decades ago, the researchers were surprised to discover inside the cells of the malaria-causing agent, *Plasmodium*, a structure that until then had only been known in plants and algae: a plastid [1],[2]. However, this one is not green, it does not contain chlorophyll. What major event in Evolution could have caused a **chloroplast** to become one of the components of *Plasmodium* cells? Let us go back to evolution: evolution: cells with a nucleus (called **eukaryotic cells**) contain simple organelles, mitochondria and chloroplasts, formed during a process of endosymbiosis called "primary" (see [Symbiosis and evolution](#)). Systematic genome analysis shows that a single primary endosymbiosis is at the origin of all "primary" chloroplasts: when a cell has "swallowed" an ancestral cyanobacterium, possibly mixed with other microbes. Figure A shows how red **algae** was built. Current biochemical and structural data indicate that the two membranes that limited the ancestral cyanobacteria were likely at the origin of the two membranes in the chloroplast envelope. Then a new endosymbiosis – called "secondary" – occurred, which saw an alga "swallowed" by a so-called secondary eukaryotic host (Figure B). A cell is then formed nesting the algae, like a Russian doll. Then the ingested alga was reduced to the point of leaving only a plastid surrounded by 4 visible membranes. The *Plasmodium* plastid – called apicoplast – is the result of one of these events, and molecular data show that it is derived from red algae. Endosymbiosis has benefited both nested cells and has been maintained throughout the evolution. To the point that the metabolic links within the ancestor of *Plasmodium* have been consolidated and that this association has become inseparable. To conclude this marriage, important regions of DNA from the nucleus and plastid of the red algae migrated to the nucleus of the secondary host. As a result of this process of cellular inclusion and DNA incorporation, *Plasmodium* contains in its nuclear genome hundreds of genes that code for proteins originating from the red algae initially ingested. As for the apicoplast, it is a fossil relic of this red algae.

References and notes

Cover image. [Source: © NIAID (Plasmodium falciparum parasite) (CC BY 2.0) via Wikimedia Commons; source Figures A & B: © Eric Maréchal]

[Registering](#)

[1] McFadden GI, Reith ME, Munholland J & Lang-Unnasch N. (1996) Plastid in human parasites. *Nature* 381, 482.

[2] Kohler S, Delwiche CF, Denny PW, *et al* (1997) A plastid of probable green algal origin in Apicomplexan parasites. *Science* 275, 1485-1489.

[3] Biot C, Botté CY, Dubar F & Maréchal E. (2012) Malaria: Research into new therapeutic treatments targeting the apicoplast, a cellular organelle of algal origin. *Med Sci.* 28: 163-171.

L'Encyclopédie de l'environnement est publiée par l'Université Grenoble Alpes.

Les articles de l'Encyclopédie de l'environnement sont mis à disposition selon les termes de la licence Creative Commons Attribution - Pas d'Utilisation Commerciale - Pas de Modification 4.0 International.
